

The serendipitous discovery of a new green chemistry method

Dr Petri Turhanen was working on a synthesis of a modified version of the biological molecule adenosine triphosphate (ATP), when he discovered that the cation exchange resin he was using was unintentionally producing another molecule. Closer investigation revealed that the molecule had been iodinated, meaning an iodine atom had been added. This specific reaction is challenging to perform using current synthesis techniques, which often rely on toxic and hazardous chemicals and rough reaction conditions.



As an environmental crisis looms, reducing our impact on the planet becomes more pertinent than ever. You may find reassurance in making conscientious lifestyle choices, but for many there's a problem which is hard to ignore: what about the less visible use of resources? What about the waste and the transportation that occur at the industrial level – those processes which provide us with medicines, cosmetics and other products?

Managing the environmental impact of these processes often falls under the remit of 'green chemistry', the concept of designing chemical processes with minimal impact on the environment. The demand for technological solutions is increasing, and the hope is that we humans can reduce our dependence on harmful and non-renewable resources, and limit hazardous by-products of industry, while maintaining or improving on our current quality of life.

CONNECTING THE CARBONS

Much of organic chemistry – building carbon-based molecules – concerns the creation of bonds between carbon and other atoms, such as nitrogen, oxygen, boron, sulfur, phosphorus or a second carbon. Connecting these atoms is the holy grail of organic chemistry because it allows chemists to build up different kinds of organic compounds. Because life is 'carbon-based', many of the most important medicines are also carbon-based, organic molecules.

But creating these bonds isn't always simple. Molecules containing carbon-


iodine (C-I) bonds are often used as intermediate molecules, like a scaffolding which makes synthesising bigger organic molecules possible. This is because not only are C-I bonds relatively easy to form, but iodine is also fairly easy to replace with another atom.

These 'iodinated' compounds can also be used for more than just simplifying synthesis. Iodine has important isomers, like ^{123}I , ^{124}I and ^{125}I , which are radioactive. Radioactive iodinated organic molecules are used in health applications like positron emission tomography (PET scanning), cancer therapy and diagnostics. But the chemical reactions performed at an industrial level to create these important iodinated compounds often use toxic starting materials, harmful solvents and even heavy metals as catalysts.

METHODS OF IODINATING

The traditional methods for iodinating molecules may use I_2 as a source of iodide, and for some reactions this works very well. However, for reactions which require a higher temperature, there is a problem. With very little heat, iodine can sublime, meaning it passes straight from the solid state to being a gas.

For some compounds, other methods are needed to successfully iodinate. But these methods could hardly be considered environmentally friendly: they involve hazardous reactants like hydrogen peroxide, trimethylsulfonium iodate and iodine monochloride, and solvents like dichloromethane, and dioxane and tetrahydrofuran – compounds which are harmful, if not



toxic, carcinogenic, environmentally damaging or explosive.

UNEXPECTED DISCOVERY

This is where Dr Petri Turhanen, researcher at the University of Eastern Finland, made an unexpected discovery. While developing a synthesis route for a derivative of ATP, an important molecule in cells, his research group found that amongst their intended product was another product they weren't expecting. Running NMR tests on this compound confirmed its identity. Where the starting material had a double bond between two carbon atoms, this molecule had a single bond, and an extra atom on both carbons – one hydrogen and one iodine.

ADDITION REACTIONS

This type of reaction – breaking a double carbon bond to add two atoms – is known as an addition reaction. There are a few methods to perform addition of hydrogen and iodine across carbon double bonds, provided the double bond is part of a longer chain. But this specific addition reaction, of hydrogen and iodine across a carbon double bond at the end of a chain, is extremely uncommon in the literature. So why had this synthesis never been found before?

Part of the answer lies in the specific materials being used to complete this reaction. Dr Turhanen's group were using a substance known to many as Dowex®, a solid resin made from the chemical polystyrene sulfonate. Dowex® is a cation exchange resin,

meaning the resin 'stores' positive ions, which it can swap for other positive ions. An example of the use of this material is in water softening, where hard water is passed through the sodium form of Dowex®. The resin absorbs the calcium and magnesium ions, releasing sodium ions into the water. And because Dowex®'s uses

They have tested the iodides of potassium, rubidium, nickel, ammonium, manganese and antimony, but the best yields are typically achieved using the readily available sodium iodide.

INITIAL TESTING

Initially, Turhanen and his team tested over 50 reaction types, finding many

Many of the compounds produced have only been reported a few times and a couple of them were previously completely unknown.


are so specific, this is the first time it had been used to perform this kind of addition reaction.

CHOOSING THE BEST IODIDE SOURCE

Since discovering this unconventional use of Dowex®, Turhanen and his team have been systematically investigating as many variations on this reaction as possible.

new methods to produce iodine-containing molecules which extend far beyond addition of hydrogen and iodine across a carbon-carbon double bond.

One of the first reactions which Turhanen discovered was the ring opening reaction of cyclic ethers (rings of carbon where one atom is an oxygen), which was previously



Dowex® H⁺ cation exchange resin before (left) and after (right) drying in oven at 120 °C overnight.

impossible for some specific cyclic ether molecules. This reaction creates molecules known as iodoalkanols, which contain an alcohol (OH) group at one end and an iodine atom at the other. Iodoalkanols are valuable molecules in organic synthesis, but are commonly made using reagents, which can be environmentally damaging. Using a cation-exchange resin and sodium iodide, Turhanen's method is much more environmentally friendly. Some cyclic ethers are particularly difficult to open, like 1,4-dioxane, opening of which has not been reported until now. Most of the reactions so far require toxic substances such as barium trifluoride. This is no longer necessary using the new method.

DI-IODIDES

More recently, Turhanen has reported the ability of Dowex® and sodium iodide to convert ethynes – molecules with a carbon-carbon triple bond – to di-iodides, adding an iodine atom to both ethyne carbon atoms. Testing of 12 readily available ethyne molecules showed that this process is versatile and provides yields of up to 78%. Many of the compounds produced have only been reported a few times before in the literature, and two of them were previously completely unknown.

ESTERIFICATION

While playing with the parameters for these reactions, Turhanen and his research group stumbled upon another neat discovery: while using alcohols as a solvent with carboxylic acids, some more unexpectedly mild reaction conditions to create esters were found. Esters are compounds which make a taste, for example, apple. These reactions are by no means new in chemistry, but they are important areas of development in green chemistry because the methyl esters of fatty acids are the main component of biodiesel. Esterification typically occurs using temperatures of up to 100 °C over the course of several hours or even days. Using Dowex®, Turhanen has been able to esterify some free fatty acids at room temperature in just 15 minutes, with yields of 99%. Extending this to a more realistic scenario for creating biodiesel, Turhanen took samples of used cooking



Chemists can increasingly choose 'greener' methods for organic synthesis.

oil from local restaurants and converted them into biodiesel, with purity greater than 95%.

A similar reaction allows the group to create esters from alcohols and carboxylic acids in mild conditions. Natural amino acids can be esterified without amino group protection. Certain types of ethers can also be prepared by the method.

One of the most exciting applications of this is selective etherification of one of the alcohol groups in glucose. The traditional method for performing this reaction is a multi-step process using bulky 'protecting' groups which only allow the desired part of the glucose molecule to react. Using Turhanen's method, the entire reaction can occur as a 'one-pot', with limited steps, lower waste and limited reagents.

GREEN CHEMISTRY


The property of cation exchange resins used alongside sodium iodide to

function as a reagent or catalyst in so many reactions makes this a valuable tool for organic synthesis, particularly because some of the reactions have never previously been possible.

One of the most exciting aspects of this discovery is the impact it will have on how some organic chemicals are synthesised. In an era where

green chemistry is a necessary part of sustaining global welfare, this method can viably replace many of the more environmentally damaging reaction conditions. Where harmful reagents, solvents and catalysts have traditionally been used, organic chemists can now opt for the simple, green solvent 2-propanol; instead of multi-step complex reactions, some synthetic routes are now possible using a 'one pot' method; and where toxic heavy metal catalysts are currently used, a reusable resin can be just as effective even at lower temperatures.

Using Turhanen's method, the entire reaction can occur as a 'one-pot', with limited steps, lower waste and limited reagents.



Behind the Research

Dr Petri Turhanen

E: petri.turhanen@uef.fi **in** <https://www.linkedin.com/in/petri-turhanen-1b67985/>
W: https://www.researchgate.net/profile/Petri_Turhanen

Research Objectives

Dr Turhanen works in green chemistry, specifically developing new techniques to produce chemicals while minimising environmental impacts.

Detail

Dr Petri Turhanen
University of Eastern Finland
Kuopio Campus
School of Pharmacy
Yliopistonranta 1 C, Mediteknia
Room 3056
P.O. BOX 1627
FI-70211 Kuopio, Finland

Bio

Dr Petri Turhanen is a university researcher at University of Eastern Finland (UEF), School of Pharmacy. He received his PhD in synthetic organic and medicinal chemistry at University of Kuopio in 2003. He has over 20 years' experience in synthetic chemistry.

Funding

Research has been supported by the Jane and Aatos Erkko Foundation.

Collaborators

- Prof Jouko J. Vepsäläinen
- Dr Juri M. Timonen and Dr Jukka Leppänen



UNIVERSITY OF
EASTERN FINLAND

References

Timonen, J., & Turhanen, P. (2019). (E)-Di-iodination of Alkynes Using Dried Dowex H+/NaI Approach. *ACS Omega*, 4(11), 14663-14668.

Turhanen, P., Leppänen, J., & Vepsäläinen, J. (2019). Green and Efficient Esterification Method Using Dried Dowex H+/NaI Approach. *ACS Omega*, 4(5), 8974-8984.

Turhanen, P., & Vepsäläinen, J. (2016). Preparation of useful building blocks, α -iodo- and bromoalkanols from cyclic ethers using the Dowex H+/NaX (X = I, Br) approach. *RSC Advances*, 6(19), 15937-15940.

Turhanen, P., & Vepsäläinen, J. (2015). A powerful tool for acid catalyzed organic addition and substitution reactions. *RSC Advances*, 5(33), 26218-26222.

Personal Response

Of the potential applications of these new reaction conditions, which do you find most exciting?

First, the most exciting application was the ease of synthesis of biodiesel from used cooking oil. Actually, according to my knowledge the method we found is the only one which also works for partially hydrolyzed cooking oil without forming soap as current industrial methods used to produce biodiesel do. Second, Dowex® is reusable in most cases such as esterifications and does not even need re-activation or regeneration, which makes the method 'green'. Third, using the method opens novel reactions, like those we have shown, for example, opening for the first time the 1,4-dioxane to $I(CH_2)_2O(CH_2)_2OH$ which is an important precursor of polyglycol synthesis. And there are more to come...